

DNA Topoisomerases

Chemistry & Biology
Review

17, May 28, 2010



DNA Topoisomerases and Their Poisoning by Anticancer and Antibacterial Drugs

Yves Pommier,^{1,*} Elisabetta Leo,¹ HongLiang Zhang,¹ and Christophe Marchand¹

¹Laboratory of Molecular Pharmacology, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD 20892-4255, USA

*Correspondence: pommier@nih.gov



Reviews

pubs.acs.org/acschemicalbiology

2013

Drugging Topoisomerases: Lessons and Challenges

Yves Pommier*

Laboratory of Molecular Pharmacology, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, Maryland, United States

Review

REVIEWS

NATURE REVIEWS | MOLECULAR CELL BIOLOGY

ADVANCE ONLINE PUBLICATION


October 2016

Roles of eukaryotic topoisomerases in transcription, replication and genomic stability

Yves Pommier¹, Yilun Sun², Shar-yin N. Huang¹ and John L. Nitiss²


Abstract | Topoisomerases introduce transient DNA breaks to relax supercoiled DNA, remove catenanes and enable chromosome segregation. Human cells encode six topoisomerases (TOP1, TOP1mt, TOP2 α , TOP2 β , TOP3 α and TOP3 β), which act on a broad range of DNA and RNA substrates at the nuclear and mitochondrial genomes. Their catalytic intermediates, the topoisomerase cleavage complexes (TOPcc), are therapeutic targets of various anticancer drugs. TOPcc can also form on damaged DNA during replication and transcription, and engage specific repair pathways, such as those mediated by tyrosyl-DNA phosphodiesterase 1 (TDP1) and TDP2 and by endonucleases (MRE11, XPF–ERCC1 and MUS81). Here, we review the roles of topoisomerases in mediating chromatin dynamics, transcription, replication, DNA damage repair and genomic stability, and discuss how deregulation of topoisomerases can cause neurodegenerative diseases, immune disorders and cancer.

DNA Topoisomerases And Cancer




Topoisomerases and TOP Genes in Humans

Not counting SPO11, there are 3 types of topoisomerases and 6 TOP genes in humans




Humans vs. Escherichia Coli

Humans have 3 types of topoisomerases and 6 TOP genes while Escherichia Coli has 2 types of topoisomerases and 6 genes



Topoisomerase differences

Biochemical differences between Top1 and Top2



Comparisons

Comparison of the 6 human topoisomerases


Genes	Chromosome	Proteins	Localization	Drugs	Mechanism	Polarity*	Main functions
<i>TOP1</i>	20q12-q13.1	Top1 100 kDa monomer	Nucleus	Camptothecins Iridinoids (LMPS)	Swivelling controlled rotation dsDNA	3'-PY	Nuclear supercoiling relaxation
<i>TOP1MT</i>	8q24.3	Top1mt 100 kDa monomer	Mitochondria	none			mitochondrial supercoiling relaxation
<i>TOP2A</i>	17q21-q22	Top2α 170 kDa dimer	Nucleus Mitochondria	Anthracyclines, (doxorubicin) Etoposide mitoxantrone	Strand passage dsDNA ATPase	5'-PY	Decatenation/replication
<i>TOP2B</i>	3p24	Top2β 180 kDa dimer	Nucleus Mitochondria				Transcription; Unknotting
<i>TOP3A</i>	17p12-p11.2	Top3α 100 kDa monomer	Nucleus Mitochondria	none	Strand passage within single strands	5'-PY	DNA Replication with BLM**
<i>TOP3B</i>	22q11.22	Top3β 100 kDa monomer	Nucleus cytoplasm				RNA topoisomerase with TDRD3

*: Covalent linkage between the catalytic tyrosine and the end of the broken DNA

** : Bloom syndrome, RecQ helicase

Topoisomerase and genomes

Topoisomerases and tyrosyl DNA phosphodiesterases (TDPs) handle both the nuclear and mitochondrial genomes and their imbalance is source of genomic instability



Top 1


Top1

TOP1 (nuclear Top1)

TOP1MT (mitochondrial Top1)




Top1 and Top2 differences




Relaxation of DNA

Relaxation of DNA by Topoisomerase I (top1)



Top1 is essential for transcription and replication (repair?)

Top1



DNA supercoiling

DNA supercoiling


In the context of chromatin, where the rotation of DNA is constrained, DNA supercoiling (over- and under-twisting and writhe) is readily generated. TOP1 and TOP1mt remove supercoiling by DNA untwisting, acting as “swivelases”, whereas TOP2 α and TOP2 β remove writhe, acting as “writhases” at DNA crossovers (see TOP2 section).

Here are some basic facts concerning DNA supercoiling that are relevant to topoisomerase activity:

- Positive supercoiling (Sc+) tightens the DNA helix whereas negative supercoiling (Sc-) facilitates the opening of the duplex and the generation of single-stranded segments.
- Nucleosome formation and disassembly absorbs and releases Sc-, respectively.
- Polymerases generate Sc+ ahead and Sc- behind their tracks.
- Excess of Sc+ arrests DNA tracking enzymes (helicases and polymerases), suppresses transcription elongation and initiation, and destabilizes nucleosomes.
- Sc- facilitates DNA melting during the initiation of replication and transcription, D-loop formation and homologous recombination and nucleosome formation.
- Excess of Sc- favors the formation of alternative DNA structures (R-loops, guanine quadruplexes, right-handed DNA (Z-DNA), plectonemic structures), which then absorb Sc- upon their formation and attract regulatory proteins.


The Two Human Top 1s

The two human Top1s





Camptothecin

Camptothecin and its derivatives used for the treatment of cancers




Camptothecin is an alkaloid from *Camptotheca acuminata* Decne, a rapidly growing tree from China. Discovered by Monroe Wall and Mansukh Wani who also discovered taxol.



Interfacial inhibitor


Camptothecins as one of Nature's Paradigms for Interfacial Inhibitors



Mutation

**Top1 is the only cellular target for camptothecin
=> Camptothecins are highly Targeted Therapies**

- **Camptothecin-producing plants encode N722S mutation (Saito and coworkers, PNAS 2008)**
- **The N722S mutation was first found in human leukemia CEM cells selected for CPT resistance (Cancer Res 2001)**



Why New Top1 Inhibitors?

Why New Top1 Inhibitors?


1. Because camptothecins are effective anticancer drugs.
Hence, Top1 is a validated target for cancer treatment.
2. Because agent with a common target have different pharmacology, toxicology and exhibit different anticancer activity (for instance top2 poisons or tubulin inhibitors [colchicine <-> vinblastine]).
3. Because camptothecins have limitations:
 - Bone marrow and intestinal toxicity (adults).
 - Drug efflux substrates (ABCG2).
 - Chemically unstable: E-ring opening.



Pharmacological Limitations of Camptothecins:

Pharmacological limitations of camptothecins:

1. Unstable at physiological pH



2. Camptothecins bind reversibly to the top1 cleavage complexes. Hence cleavage complexes reverse rapidly after drug removal => prolonged infusions


Indenoisoquinolines

Clinical development of the indenoisoquinolines

Indenoisoquinolines

Two compounds selected LMP-776 and LMP-400:

- Potent and specific Top1cc-targeted drugs
- Overcoming limitations of camptothecins:
 - Chemical stability (no alpha-hydroxylactone)
 - Overcome resistance drug efflux pumps (collaboration with S. Bates & M. Gottesman)
- Trap Top1cc at different genomic sites compared to camptothecins => target $\neq$ regions of the genome.



Targeted delivery

Second Generation Camptothecins with Targeted Delivery

Name	Company	Active Derivative (Payload)	Formulation (Conjugate; Target)
Onivyde™ = MM398*	Merrimack	Irinotecan	Liposome
CRLX101	Cerulean Pharma Inc.	Camptothecin	PEG
NKTR-102	Nektar Therapeutics	Etirinotecan (20-position)	PEG (Pegol)
IMMU-132	Immunomedics	SN-38 (20-position)	CDA - TROP2 (TACSD2)
STA-12-8666	Synta Pharmaceuticals	SN-38 (10-position)	HDC – Hsp90

** FDA approved October 2015*

Comparative oncology trials

Comparative Oncology Trials Consortium

CCR-COP website

Create

The C
Reser
biolog
pharm
institu

All vet



For Staff | Login

Search CCR...

Search

Cancer
er
id
r

[COP HOME](#)

[PET OWNERS](#)

[RESEARCH](#)

[TRIAL SPONSORS](#)

[Home](#) > Comparative Oncology Program

COP | comparative oncology program



Goals:

1. Compare LMP400, LMP776 and LMP744
2. Determine MTD in dogs with lymphomas
3. Determine and compare activity of 3 drugs
4. Determine pharmacokinetics in blood and tumor
5. Determine target engagement:
 1. γ H2HAX
 2. TOP1 downregulation

[Pet Owners](#)


[Research](#)

[Trial Sponsors](#)

[Clinical Trials](#)

Dog lymphoma

All drugs exhibit antitumor activity in primary dog lymphoma



Amy LeBlanc
CCR COP



James Doroshow
DCTD - CCR

Indotecan and imidotecan trials

Summary of the clinical oncology trial:

- The two clinical indenoisoquinolines, LMP400 (indotecan) and LMP776 (imidotecan) exhibit antitumor activity in dog lymphoma.
- The 3rd indenoisoquinoline, LMP744 shows even greater antitumor activity.
- The dose limiting toxicity of the indenoisoquinolines (MTD = 17.5 mg/m² for LMP776; MTD > 65 mg/m² for LMP400; MTD = 100 mg/m² for LMP744) is bone marrow suppression. No diarrhea.
- The PK of the LMPs shows long half-lives: LMP744: 17 h; LMP400: 11 h; LMP776: 6 h.
- LMP744 shows remarkable tumor retention and accumulation
- γH2AX response demonstrates target engagement for all drugs

Precision therapeutics

Precision therapeutics can be defined as the ability to:

1. prescribing effective therapies only to those patients who will respond effectively (cure) ⇔ **Tumor molecular signature**: SLFN11 + HRD...
2. while limiting toxicity to normal tissues and minimizing side effects
⇔ **Targeted delivery**



Camptothecins

Second Generation Camptothecins with Targeted Delivery

Name	Company	Active Derivative (Payload)	Formulation (Conjugate; Target)
Onivyde™ = MM398*	Merrimack	Irinotecan (CPT11)	Liposome
CRLX101	Cerulean Pharma Inc.	Camptothecin	PEG
NKTR-102	Nektar Therapeutics	<u>Etirinotecan</u> (20 position)	PEG (<u>Pegol</u>)
PLX038	ProLynx	SN-38	PEG
IMMU-132 = <u>Sacituzumab</u> <u>govitecan</u>	Immunomedics (Seattle Genetics)	SN-38 (20 position)	ADC - TROP2 (TACSD2)
IMMU-130 = <u>Labetuzumab</u> <u>govitecan</u>	Immunomedics	SN-38	ADC-CEACAM5
DS-8201a ***	Daichi Sankyo	<u>DXd</u> (<u>Exatecan</u>)	ADC - HER2
PEN-866	<u>Tarveda</u> Therapeutics	SN-38 (10 position)	HDC - Conjugate Hsp90
NK012	Nippon Kayaku	SN-38	Polymeric micelles (PEG-polyglutamate)
ALOS4-CPT	Ariel University	Camptothecin	HDC - ALOS-4
SN38-TOA	CHOP Philadelphia	SN-38	Tocopherol <u>oxyacetate</u> nanoparticles

* FDA Approved, October 2015

** FDA Breakthrough, February 2016

*** FDA Breakthrough, August 2017 (Breast)

Camptothecins as
warheads

Tumor-specific delivery

Top2

Top2

Top2 α – TOP2A: Replication




Highly expressed in replicating and cancer cells

Top2 β – TOP2B: Transcription

Expressed both in replicating and differentiated cells




Two Top2 enzymes



Humans have two Top2 enzymes

Top2

Top2 catalyze a broad range of reactions



Transcription

Pommier, Y., Sun, Y., Huang, S. & Nitiss, J.L.
2016 Nature Rev Mol Cell Biol





Figure 2 | Topoisomerases and transcription. Transcription incurs topological constraints that result from the progression of RNA polymerase II (Pol II). Positive supercoiling (Sc⁺) of the DNA template takes place ahead of the transcription bubble, which in turn obstructs further Pol II movement, and negative supercoiling (Sc⁻), which promotes the formation of RNA–DNA hybrids (R loops), accumulates behind it. TOP2 and especially TOP1 enzymes function ahead of Pol II to remove positive supercoils, whereas relaxation of negative supercoils behind the transcription apparatus relies on TOP1 and TOP3β. In addition, TOP1 regulates the activity of the transcription factor TATA-box-binding protein (TBP) at promoter TATA boxes independently of its catalytic activity. The formation of TOP2β-mediated transient DNA double-stranded breaks at promoter regions in certain genes is crucial for transcription activation. TOP1 is also recruited to certain enhancer regions to promote (ligand-dependent) enhancer activation by generating transient DNA single-stranded breaks. Topological barriers are genomic regions where the DNA is not free to rotate around its axis and require TOP1 and TOP2 to relax supercoils (Sc). TF, transcription factor.


DNA replication




Functions of topoisomerases in DNA replication. **a.** Initiation of DNA replication requires separation of the two parental strands, which generates negative supercoiling (Sc⁻) at the origin of replication and positive supercoiling in the flanking regions due to topological barriers, such as nuclear matrix attachment sites or insulators. Positive supercoiling is dissipated by TOP1 and TOP2α to allow replication fork progression (arrows). **b.** Replication elongation generates positive supercoiling ahead of the replication fork and negative supercoiling behind it. Positive supercoiling is removed by TOP1 and TOP2α, whereas negative supercoiling can be removed by TOP1, TOP2α or TOP3α. TOP2α can also remove precatenanes, which are formed when the fork rotates during elongation. **c.** Converging forks generate high positive supercoiling between them. **d.** Upon replication completion, catenanes are removed by TOP2α (left) and hemicatenanes by TOP3α (right). Topological barriers are genomic regions where the DNA is not free to rotate around its axis, for example owing to hindrance by macromolecular complexes.

Top2 drugs


Anticancer Top2-targeted drugs



Antibiotics Top2-targeted drugs




Etoposide



Structure of a
topoisomerase II
cleavage complex
(Top2cc) trapped
by etoposide (VP-
16)

Levofloxacin

Antibacterials



Structure of a topoisomerase IV cleavage complex (Topo IVcc) trapped by the quinolone, levofloxacin

Interfacial inhibition

TRENDS in Pharmacological Sciences Vol.26 No.3 March 2005



Review

TRENDS in Pharmacological Sciences Vol.26 No.3 March 2005

Full text provided by www.sciencedirect.com



Interfacial inhibition of macromolecular interactions: nature's paradigm for drug discovery

Yves Pommier¹ and Jacqueline Cherfils²

¹Laboratory of Molecular Pharmacology, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD 20892-4255, USA

²Laboratoire d'Enzymologie et Biochimie Structurales, CNRS, Gif sur Yvette, France

NATURE REVIEWS | **DRUG DISCOVERY**


VOLUME 11 | JANUARY 2012

Interfacial inhibitors: targeting macromolecular complexes


Yves Pommier¹ and Christophe Marchand¹

Abstract | Interfacial inhibitors belong to a broad class of natural products and synthetic drugs that are commonly used to treat cancers as well as bacterial and HIV infections. They bind selectively to interfaces as macromolecular machines assemble and are set in motion. The bound drugs transiently arrest the targeted molecular machines, which can initiate allosteric effects, or desynchronize macromolecular machines that normally function in concert. Here, we review five archetypical examples of interfacial inhibitors: the camptothecins, etoposide, the quinolone antibiotics, the vinca alkaloids and the novel anti-HIV inhibitor raltegravir. We discuss the common and diverging elements between interfacial and allosteric inhibitors and give a perspective for the rationale and methods used to discover novel interfacial inhibitors.

Topoisomerase drugs



Camptothecins
Indenoisoquinolines



Etoposide
Doxorubicin

Top 3

Top3

Top3 α – TOP3A: Replication

DNA topoisomerase (single-strands);
resolves hemicatenanes and
prevents recombinations


Top3 β – TOP3B: Transcription

DNA topoisomerase (R-loops);
RNA topoisomerase



Decatenation

Decatenation Top2 vs. Top3



Top-3beta

Nature NeuroScience 2013

Deletion of TOP3 β , a component of FMRP-containing mRNPs, contributes to neurodevelopmental disorders

Georg Stoll^{1,32}, Olli P H Pietiläinen^{2-4,32}, Bastian Linder^{1,32}, Jaana Suvisaari⁵, Cornelia Brosi¹, William Hennah^{3,5}, Virpi Leppä³, Minna Torniainen⁵, Samuli Ripatti^{2,3}, Sirpa Ala-Mello⁶, Oliver Plöttner⁷, Karola Rehnström², Annamari Tuulio-Henriksson⁵, Teppo Varilo^{3,4}, Jonna Tallila², Kati Kristiansson³, Matti Isohanni⁸, Jaakko Kaprio^{3,5,9}, Johan G Eriksson¹⁰⁻¹⁴, Olli T Raitakari^{15,16}, Terho Lehtimäki¹⁷, Marjo-Riitta Jarvelin¹⁸⁻²¹, Veikko Salomaa²², Matthew Hurles², Hreinn Stefansson²³, Leena Peltonen^{2-4,24,25}, Patrick F Sullivan^{26,27}, Tiina Paunio^{3,4,28}, Jouko Lönnqvist^{3,6}, Mark J Daly^{29,30}, Utz Fischer¹, Nelson B Freimer³¹ & Aarno Palotie^{2,3,30}

Implicating particular genes in the generation of complex brain and behavior phenotypes requires multiple lines of evidence. The rarity of most high-impact genetic variants typically precludes the possibility of accruing statistical evidence that they are associated with a given trait. We found that the enrichment of a rare chromosome 22q11.22 deletion in a recently expanded Northern Finnish sub-isolate enabled the detection of association between TOP3B and both schizophrenia and cognitive impairment. Biochemical analysis of TOP3 β revealed that this topoisomerase was a component of cytosolic messenger ribonucleoproteins (mRNPs) and was catalytically active on RNA. The recruitment of TOP3 β to mRNPs was independent of RNA cis-elements and was coupled to the co-recruitment of FMRP, the disease gene product in fragile X mental retardation syndrome. Our results indicate a previously unknown role for TOP3 β in mRNA metabolism and suggest that it is involved in neurodevelopmental disorders.

Top3 β is an RNA topoisomerase that works with fragile X syndrome protein to promote synapse formation


Dongyi Xu^{1,2,10}, Weiping Shen^{1,10}, Rong Guo¹, Yutong Xue¹, Wei Peng¹, Jian Sima³, Jay Yang⁴, Alexei Sharov⁵, Subramanya Srikantan⁶, Jiandong Yang¹, David Fox III¹, Yong Qian⁵, Jennifer L Martindale⁶, Yulan Piao⁵, James Machamer⁷, Samit R Joshi⁸, Subhasis Mohanty⁸, Albert C Shaw⁸, Thomas E Lloyd⁷, Grant W Brown⁴, Minoru S H Ko⁵, Myriam Gorospe⁶, Sige Zou⁹ & Weidong Wang¹

Topoisomerases are crucial for solving DNA topological problems, but they have not been linked to RNA metabolism. Here we show that human topoisomerase 3 β (Top3 β) is an RNA topoisomerase that biochemically and genetically interacts with FMRP, a protein that is deficient in fragile X syndrome and is known to regulate the translation of mRNAs that are important for neuronal function, abnormalities of which are linked to autism. Notably, the FMRP-Top3 β interaction is abolished by a disease-associated mutation of FMRP, suggesting that Top3 β may contribute to the pathogenesis of mental disorders. Top3 β binds multiple mRNAs encoded by genes with neuronal functions linked to schizophrenia and autism. Expression of one such gene, that encoding protein tyrosine kinase 2 (ptk2, also known as focal adhesion kinase or FAK), is reduced in the neuromuscular junctions of Top3 β mutant flies. Synapse formation is defective in Top3 β mutant flies and mice, as well as in FMRP mutant flies and mice. Our findings suggest that Top3 β acts as an RNA topoisomerase and works with FMRP to promote the expression of mRNAs that are crucial for neurodevelopment and mental health.


Top3A and Top3B

TOP3 alpha and beta function in different protein complexes and biological processes

Replication



Transcription



Topoisomerases

Topoisomerases
Genomic Integrity
and
Human diseases



DNA damage

Topoisomerase-induced DNA damage

A



- * Conversion of TOP1cc into DSB by replication "run-off"
=> TOP1 needs to be removed by TDP1
and/or 3'-flap endonucleases (XPF-ERCC1)
=> DSB repaired by homologous recombination
- * TOP1cc also form DSB when on opposite strands or
opposite to a preexisting single-strand break

B



- * TOP2cc readily form DSB when concerted cleavage
on both strands and disjunction of the homodimer

C



- * Collisions of polymerases and helicases (green ellipse)
with trapped Top cleavage complexes (Stop sign)
=> Protein-DNA complexes blocking DNA metabolism

D



- * Topological defects resulting from enzyme sequestration
in the cleavage complexes: accumulation of
=> supercoils (Top1 and Top2) (1)
=> knots (Top2) (2)
=> catenanes (Top2) (3)

Topoisomerases and disease


Table 1 Drugs, DNA alterations and physiological processes that lead to the formation of persistent TOPcc		
Causes	Consequences for TOP1 enzymes	Consequences for TOP2 enzymes
Anticancer drugs acting as interfacial inhibitors ¹⁵⁵	Trapping of TOP1cc by irinotecan, topotecan, indenoisoquinolines* and tumour-targeting camptothecin derivatives ^{3,154,155}	Trapping of TOP2cc by etoposide, teniposide, doxorubicin, epirubicin, idarubicin and mitoxantrone ⁴
Oxidative DNA lesions (8-oxoguanine, 8-oxoadenosine and 5-hydroxycytosine)	Induction and trapping of TOP1cc ^{218,219}	Induction and trapping of TOP2cc ²²⁰
Abasic sites and DNA mismatches	Formation of irreversible TOP1cc ²²¹	Formation of irreversible TOP2cc ^{220,222–225}
Carcinogenic base adducts (methylated bases, exocyclic adducts, benzo[a]pyrene adducts and crotonaldehyde adducts)	Induction and trapping of TOP1cc ^{226–232}	Induction and trapping of TOP2cc ^{220,233–235}
Nicks and DNA strand breaks	Formation of irreversible TOP1cc, double-stranded breaks, genomic deletions and recombination ^{18,167,168,236,237}	Formation of irreversible TOP2cc ²³⁵
UV lesions (pyrimidine dimers and 6.4-photoproducts)	Induction of TOP1cc ^{238,239}	Enzymatic inhibition ²⁴⁰
Ribonucleotide incorporation into DNA	Formation of TOP1cc that generate nicks with 2',3'-cyclic phosphate ends and short deletions in repeat sequences ^{166–168}	Stabilization of TOP2cc with asymmetrical cleavage ^{20,169,241}
Natural and food products	Unknown	Stabilization of TOP2cc by flavones, tea and wine products ²⁰⁵
Genetic defects	Unrepaired TOP1cc due to TDP1 defects ^{177,206,210} in cooperation with ATM defects ¹⁷⁹	Unrepaired TOP2cc due to TDP2 defects ⁶⁹
Transcription activation	Stabilization of TOP1cc at enhancers ⁴²	Stabilization of TOP2cc at promoters ^{62,65,242,243}

ATM, ataxia telangiectasia mutated; TDP, tyrosyl-DNA phosphodiesterase; TOPcc, topoisomerase cleavage complex.

*Indenoisoquinoline derivatives are in clinical trials.

Replicative DNA damage

Replicative DNA damage induced by TOP1cc (Topoisomerase I cleavage complexes)



Human diseases

Human Diseases linked with topoisomerases

TOP1: Neurological diseases due to lack of removal of TOP1cc (in conjunction with TDP1 and ATM deficiencies)

TOP2B: Chromosome translocations at TOP2Bcc (leukemia, prostate cancers...)

TOP3B: Neurodevelopmental disorders (schizophrenia and cognitive impairment)

TDP1: SCAN1 (Spinocerebellar Ataxia and peripheral Neuropathy)

TDP2: Intellectual disability, seizures and ataxia

DNA repair

Box 1 | DNA–protein crosslink repair pathways and human health

It is intriguing that germline mutations in almost all identified genes that encode components of the three main DNA–protein crosslink (DPC) repair pathways result in human syndromes that are characterized by genome instability, cancer predisposition, premature ageing and/or neurological pathologies. Whether all of these phenotypes are directly related to a defect in DPC repair or to other cellular functions of these proteins, is not entirely clear in all cases. The MRN complex, for example, has crucial functions during repair of DSBs, which are clearly related to the radiosensitivity and immunodeficiency that are observed in patients with mutations in genes that encode MRN subunits. Below, we briefly discuss the main diseases that are associated with mutations in DPC repair proteins.

Repair by tyrosyl-DNA phosphodiesterases

Spinocerebellar ataxia, autosomal recessive, with axonal neuropathy (SCAN1; OMIM: 607250) was first identified in a large Saudi Arabian family (nine affected individuals) that had homozygous mutations in the tyrosyl-DNA phosphodiesterase 1 (*TDP1*) gene, which map to chromosome 14q31–14q32 (REF. 91). Clinical features of SCAN1 include spinocerebellar ataxia (with late onset and slow progression) and areflexia, followed by signs of peripheral neuropathy, with the absence of non-neurological symptoms that are otherwise common in ataxia telangiectasia (telangiectasias, immunodeficiency, and cancer predisposition). Interestingly, the TDP1-H493R variant, which causes SCAN1, is not only catalytically compromised but also becomes covalently trapped in the process of repairing Top1 adducts⁹². However, despite this pathological gain-of-function of the TDP1-H493R variant, this form of SCAN1 is a recessive disorder, as wild-type TDP1 is able to repair the TDP1-H493R adducts in heterozygous individuals.

Spinocerebellar ataxia, autosomal recessive 23 (SCAR23; OMIM: 616949) has been identified in three Irish brothers who were born to consanguineous parents, and in an unrelated Egyptian case. SCAR23 has been associated with a homozygous mutation in the *TDP2* gene on chromosome 6p2 (REF. 40). Clinical features include progressive spinocerebellar ataxia, epilepsy and intellectual disabilities.

Repair by the MRN complex

Clinical features of ataxia telangiectasia-like disorder 1 (ATLD1; OMIM: 604391) include slowly progressive cerebellar degeneration that results in ataxia and oculomotor apraxia, and dysarthria, but without telangiectasia or major defects in immunoglobulin production, and without major cancer predisposition but with radiosensitivity. ATLD1 is caused by homozygous or compound heterozygous mutations in the *MRE11* gene on chromosome 11q21 (REFS 93,94).

Nijmegen breakage syndrome (NBS) ataxia telangiectasia variant V1 (OMIM: 251260) is caused by homozygous or compound heterozygous mutations in the *NBS1* gene on chromosome 8q21. More than 90% of patients are homozygous for a five base pair deletion (657del5), which leads to a frameshift and truncation of the NBS1 protein^{95–98}. There are no reliable estimates of worldwide prevalence, but it is likely to approximate to 1 in 100,000 live births (most common in the Slavic populations of Eastern Europe)⁹⁹. Clinical features of this syndrome include microcephaly, growth retardation, immunodeficiency, predisposition to cancer (mainly non-Hodgkin lymphoma), and radiosensitivity; neither ataxia nor telangiectasia are present. Compound heterozygous mutations in the *RAD50* gene (on chromosome 5q31.1) that give rise to low levels of RAD50 cause Nijmegen breakage syndrome-like disorder (NBSLD; OMIM 613078)¹⁰⁰. Clinical features of NBSLD include microcephaly, growth retardation, chromosome instability, radioresistant DNA synthesis, radiation hypersensitivity and slight, non-progressive ataxia; there are no signs of telangiectasia or immunodeficiency and no evidence of cancer predisposition^{100,101}.

Repair by DPC proteases

Homozygous or compound heterozygous mutations in the *SPRTN* gene (on chromosome 1q42) cause Ruijs–Aalfs syndrome (RJALS; OMIM: 616200). Clinical features of RJALS include growth retardation, early-onset hepatocellular carcinomas, micrognathia, chromosomal instability and sensitivity to genotoxic agents^{68,69}.



Covalent complexes

Repair of Topoisomerase
covalent complexes



Catalytic intermediate

*Reversible transesterifications,
differential geometry
and polarity (5' vs. 3' tyrosyl linkage)
of human topoisomerases*



All topoisomerases form a catalytic intermediate consisting of a covalent bond between one end of the break they make in DNA (and RNA for TOP3B) and their catalytic tyrosyl residue

Pommier et al. ACS Chem Rev 2009

<http://discover.nci.nih.gov/pommier/pommier.htm>

Topoisomerase




Figure 5 | TOPcc repair. a | Tyrosyl-DNA phosphodiesterase 1 (TDP1) and TDP2 (although much less efficiently and therefore shown in parentheses) cleave the TOP1 tyrosyl-DNA covalent bond (middle), releasing TOP1 and leaving a 3'-phosphate end (right) that needs to be further processed by polynucleotide kinase phosphatase (not shown). **b** | TOP2 cleavage complexes (TOP2cc) are preferentially repaired by TDP2 and much less efficiently by TDP1 (middle) in vertebrates, releasing TOP2 and leaving a 5'-phosphate (right), which can be readily ligated. Yeast, which do not encode a TDP2 orthologue, use Tdp1 to excise both Top1cc and Top2cc. In the endonuclease pathways (left), topoisomerases are released with the segment of DNA to which they are attached by the action of endonucleases: the polarity is opposite for TOP1cc (part a) and TOP2cc (part b).

Repair pathways

Parallel repair pathways for abortive topoisomerase cleavage complexes:

- Excision by two dissimilar tyrosyl DNA phosphodiesterases: TDP1 and TDP2
- Endonucleases (Mre11; NER...)



TDP1 has a broad range of DNA repair functions beyond TOP1cc repair:


- 3'-end cleansing activity: 3'-phosphoglycolates (H_2O_2 , bleomycin, IR)
- 3'-dRP (MMS, alkylating agents) (JBC)*
- Excises chain terminator nucleosides (AraC, AZT, abacavir, sapacitabine) (JBC; NAR)*; 3'-nucleosidase
- Both in the nucleus and mitochondria (EMBO J)
- Role in genomic stability in the nervous system (PNAS)*
- Coupled with PARP1 (JBC; DNAR)*
- Also excises TOP2cc (JBC)* (no TDP2 in yeast)

TDP2 also has DNA repair functions beyond TOP2cc:

- 5'-end tyrosyl-DNA phosphodiesterase: VpG unlinkase (poliovirus replication) (HPV replication)
- Crystal structures (NSMB; JBC)*: similarity with APE1 (Mg^{2+} ; 5 fingers) but different from TDP1
- Recruitment to TOP2cc by Ub (JBC)*
- Activity on TOP2cc requires denaturation/ proteolysis (JBC)*


Parallel repair pathways

Normal cells have parallel repair pathways for abortive TOP1cc



TDP1 is coupled with PARP1

Discovered this review cycle (NAR; DNAR; JBC)



PARP1 inhibitors synergize with TOP1 inhibitors

(Sci Transl Med 2014; JPET 2014)

⇒ Synthetic lethality

in Mre11- or XPF-ERCC1-deficient cancers?

Acknowledgements

Acknowledgements for Indenoisoquinolines

• **Mark Cushman** (Purdue University)

- William Bonner (LMP-CCR)
- Smitha Antony (LMP-CCR)
- George Vande Woude (DBS[®])
- Susan Bates (COB-CCR)
- Robert Wiltrout (CCR)
- Patricia Steeg (LMP-CCR)
- Lyuba Varticovski (LHC-CCR)

• **James Doroshow** (DCTD-LMP-NCI)

- Joseph Tomaszewski (DCTD-NCI)
- Jerry Collins (DTP-DCTD-NCI)
- Barbara Mroczkowski (DCTD)
- Ralph Parchment (SAIC-FCRDC)
- Bob Kinders (SAIC-FCRDC)

• **Kenneth Paull**[®] (DTP)

- Melinda Hollingshead (DTP)
- Jiuping Ji (SAIC-FCRDC)
- Joe Covey (DTP)
- Liz Glaze (DTP)
- Prabhakar Risbood (DTP)
- Jim Cradock (DTP)
- Rao Vishnuvajjala (DTP)
- Sima Hayavi (DTP)
- Tiziano DiPaolo (DCTP)
- Vali Sevastita (DTP)
- Gina Uhlenbrauck (DTP)
- Shivaani Kummar (COB-CCR)
- Giuseppe Giaccone (COB-CCR)